Symmetry transformations in Batalin-Vilkovisky formalism.

Albert Schwarz*
Department of Mathematics, University of California,
Davis, CA 95616
ASSCHWARZ@UCDAVIS.EDU

Abstract

Let us suppose that the functional S on an odd symplectic manifold satisfies the quantum master equation $\Delta_\rho e^S = 0$. We prove that in some sense every quantum observable (i.e. every function H obeying $\Delta_\rho(He^S) = 0$) determines a symmetry of the theory with the action functional S.

This short note was inspired by the paper [2] where the results of [1] were applied to obtain the description of the gauge transformations in Batalin-Vilkovisky theory. We begin with the observation that [1] contains conditions of physical equivalence of different solutions to the master equivalence and use these conditions to give a very transparent analysis of symmetry transformations in BV-approach. Let us recall some notions and results of [1].

Let us fix a manifold M provided with odd symplectic structure (P-structure). Let us suppose that the volume element in M is specified by the density ρ. We say that this volume element determines an SP-structure in M if $\Delta_\rho^2 = 0$. (Here the operator Δ_ρ acts by the formula

$$\Delta_\rho A = \frac{1}{2} div_\rho K_A, \tag{1}$$

where K_A denotes the hamiltonian vector field corresponding to A and the divergence div_ρ is calculated with respect to the density ρ.) By definition a function S satisfies the quantum master equation on an SP-manifold M if

$$\Delta_\rho e^S = 0. \tag{2}$$

Such a function S can be considered as an action functional and determines physical quantitional by means of integration over Lagrandian submanifolds of M.

*Research supported in part by NSF grant No. DMS-9201366

The following statement was proven in [1](see Lemma 4 and Eqn.(32)):
The density $\tilde{\rho} = e^{\sigma}\rho$ *determines a new* SP*-structure in* M *if and only if*

$$\Delta_{\rho}\sigma + \frac{1}{4}\{\sigma, \sigma\} = 0. \tag{3}$$

If S *is a solution to the quantum master equation(1) then* $\tilde{S} = S - \frac{1}{2}\sigma$ *satisfies the quantum master equation*

$$\Delta_{\tilde{\rho}} e^{\tilde{S}} = 0 \tag{4}$$

corresponding to the new SP*-structure. The action functional* $\tilde{S} = S - \frac{1}{2}\sigma$ *on the manifold* M *with the new* SP*-structure describes the same physics as the action functional* S *on the manifold* M *with the old* SP*-structure. In particular*

$$\int_L e^{\tilde{S}} d\tilde{\lambda} = \int_L e^S d\lambda \tag{5}$$

for every Lagrangian submanifold $L \subset M$*. (Here* $d\lambda$ *and* $d\tilde{\lambda}$ *denote the volume elements on* L *determined by new and old* SP*-structure correspondingly.)*

One can introduce the notion of quantum observable for the theory with the action S on an SP-manifold M in the following way. We will say that an even function A determines a quantum observable if

$$\Delta_{\rho}(Ae^S) = 0 \tag{6}$$

or, equivalently,if $\Delta_{\rho}A + \{A, S\} = 0$. It is easy to check that for every observable A we have also $\Delta_{\tilde{\rho}}(Ae^{\tilde{S}}) = 0$; in other words the quantum observables for the action functionals S and $\tilde{S}$ coincide. Moreover

$$\int_L Ae^{\tilde{S}} d\tilde{\lambda} = \int_L Ae^S d\lambda \tag{7}$$

for every observable A and for every Lagrangian manifold L. This equation can be considered as a little bit more precise expression of physical equivalence of action functionals S and $\tilde{S}$ than (5). The condition (6) is equivalent to the requirement that $S + \varepsilon A$ is a solution to the quantum master equation for infinitesimal ε; therefore we can derive (7) applying (4) to the functionals $S + \varepsilon A,\ \tilde{S} + \varepsilon A$ where $\varepsilon \to 0$.

Let us use the statement above in the case $\sigma = 2S$. As was mentioned in [1] the equation (2) can be represented in the form $\Delta_{\rho} e^{\sigma/2} = 0$; therefore it is satisfied for $\sigma = 2S$. We arrive to the following conclusion:

The theory with the action ρ *is physically equivalent to the theory with the trivial action functional* $\tilde{S} = 0$ *and the* SP*-structure determined by the density* $\tilde{\rho} = e^{2S}\rho$*.*

The statement above can be used to analyze the symmetry transformations in Batalin-Vilkovisky approach. If $\tilde{S} = 0$ then the symmetries can be characterized as transformations of M preserving the symplectic structure and the density $\tilde{\rho} = e^{2S}\rho$. Infinitesimal symmetry transformation correspond therefore to hamiltonian vector fields with zero divergence with respect to $\tilde{\rho}$, i.e. to functions H obeying $\Delta_{\tilde{\rho}} H = 0$. One can say therefore that infinitesimal symmetry transformations are in one-to-one correspondence with quantum observables of our theory. It is important to emphasize that the notion of observable does not change when we replace S by $\tilde{S} = 0$. Therefore we can describe symmetries also in terms of the original action S and the original measure $d\mu = \rho dx$. (This description follows immediately from the description of symmetry transformations in the formulation with $\tilde{S} = 0$; one can prove it directly using (5),(6),(7).)

We obtain that every function H satisfying $\Delta_\rho H + \{H, S\} = 0$ (every quantum observable) determines a symmetry in the following sense. Neither the action functional S, nor the density ρ are invariant with respect to an infinitesimal transformation with the Hamiltonian H, however the new action functional

$$\tilde{S} = S + \varepsilon\{H, S\} \tag{8}$$

and the new measure

$$d\tilde{\mu} = d\mu(1 + 2\varepsilon\Delta_\rho H \tag{9}$$

describe the same physics as the old action functional S and the old measure $d\mu = \rho dx$.

Here ε is an infinitesimal parameter. The formula (9) follows immediately from (1) and from the standard formula

$$d\tilde{\mu} = d\mu + \varepsilon d\mu \; div_\rho K \tag{10}$$

for the variation of measure by the infinitesimal transformation $x \to x + \varepsilon K$, where K is an arbitrary vector field.

It is essential to stress that quantum observables don't remain intact by the symmetry transformation (8),(9). Namely, one should replace a quantum observable A by the observable $\tilde{A} = A + \varepsilon\{H, A\}$.

References

1.A. Schwarz: Geometry of Batalin-Vilkovisky quantization. Commun. Math. Phys. 155,249-260 (1993)

2.A.Sen and B Zwiebach: A note on gauge transformations in Batalin-Vilkovisky theory. MIT-CTP-2240 TIFR-TH-93-38 hep-th/9309027,1993.